arXiv:1502.05913v3 [math.GN] 9 Mar 2015

STRONGLY NEAR PROXIMITY
& HYPERSPACE TOPOLOGY

J.F. PETERS$^{\alpha}$ AND C. GUADAGNI$^{\beta}$

Dedicated to the Memory of Som Naimpally

Abstract. This article introduces strongly near proximity $\stackrel{\wedge\wedge}{\delta}$, which represents a new kind of proximity called *almost proximity.* A main result in this paper is the introduction of a hit-and-miss topology on CL(X), the hyperspace of nonempty closed subsets of X, based on the strongly near proximity.

1. Introduction

Recently, we introduced *strongly far proximities* [31]. Strongly far proximities make it possible to distinguish between a weaker form of farness and a stronger one associated with the *Efremovič property.* In this article, we introduce a framework that enables us to single out when two sets are *near* and when they are *strongly near.* This article also introduces a kind of *hit and far-miss* topology on the hyperspace of all non-empty closed subsets CL(X) of a topological space X.

2. Preliminaries

Recall how a *Lodato proximity* is defined [20, 21, 22] (see, also, [27, 25]).

Definition 2.1. *Let X be a nonempty set. A Lodato proximity δ is a relation on $\mathscr{P}(X)$ which satisfies the following properties for all subsets A, B, C of X :*

P0) $A\ \delta\ B \Rightarrow B\ \delta\ A$
P1) $A\ \delta\ B \Rightarrow A \neq \varnothing$ *and* $B \neq \varnothing$
P2) $A \cap B \neq \varnothing \Rightarrow A\ \delta\ B$
P3) $A\ \delta\ (B \cup C) \Leftrightarrow A\ \delta\ B$ *or* $A\ \delta\ C$
P4) $A\ \delta\ B$ *and* $\{b\}\ \delta\ C$ *for each* $b \in B\ \Rightarrow A\ \delta\ C$

Further δ is separated , if

P5) $\{x\}\ \delta\ \{y\} \Rightarrow x = y$.

When we write $A\ \delta\ B$, we read "A is near to B", while when we write $A \not\delta B$ we read "A is far from B". A *basic proximity* is one that satisfies the Čech axioms $P0) - P3)$ [3, §2.5, p. 439]. *Lodato proximity* or *LO-proximity* is one of the simplest proximities. We can associate a topology with the space (X, δ) by considering as

---

2010 *Mathematics Subject Classification.* Primary 54E05 (Proximity); Secondary 54B20 (Hyperspaces).

Key words and phrases. Hit-and-Miss Topology, Hyperspaces, Proximity, Strongly Far, Strongly Near.

The research has been supported by the Natural Sciences & Engineering Research Council of Canada (NSERC) discovery grant 185986.

closed sets the ones that coincide with their own closure, where for a subset A we have

$$\mathrm{cl}A = \{x \in X : x\ \delta\ A\}.$$

This is possible because of the correspondence of Lodato axioms with the well-known Kuratowski closure axioms [19, Vol. 1,§4.1, 20-21].

By considering the gap between two sets in a metric space ($d(A,B) = \inf\{d(a,b) : a \in A, b \in B\}$ or ∞ if A or B is empty), Efremovič introduced a stronger proximity called *Efremovič proximity* or *EF-proximity* [16, 15].

Definition 2.2. *An* EF-proximity *is a relation on $\mathscr{P}(X)$ which satisfies $P0)$ through $P3)$ and in addition*

$$A \not\delta B \Rightarrow \exists E \subset X \text{ such that } A \not\delta E \text{ and } X \setminus E \not\delta B \text{ EF-property.}$$

A topological space has a compatible EF-proximity if and only if it is a Tychonoff space [28, §5.3] (see, also, S. Willard on Tychonoff spaces [38, §8, 52-53]).

Any proximity δ on X induces a binary relation over the powerset exp X, usually denoted as $\ll_\delta$ and named the *natural strong inclusion associated with* δ, by declaring that A is *strongly included* in B, $A \ll_\delta B$, when A is far from the complement of B, *i.e.*, $A \not\delta X \setminus B$.

By strong inclusion, the *Efremovič property* for δ can be written also as a betweenness property

(EF) If $A \ll_\delta B$, then there exists some C such that $A \ll_\delta C \ll_\delta B$.

A pivotal example of *EF-proximity* is the *metric proximity* in a metric space (X,d) defined by

$$A\ \delta\ B \Leftrightarrow d(A,B) = 0.$$

That is, A and B *either intersect or are asymptotic*: for each natural number n there is a point a_n in A and a point b_n in B such that $d(a_n, b_n) < \frac{1}{n}$.

2.1. **Hit and far-miss topologies.**

Let $\mathrm{CL}(X)$ be the hyperspace of all non-empty closed subsets of a space X. *Hit and miss* and *hit and far-miss* topologies on $\mathrm{CL}(X)$ are obtained by the join of two halves. Well-known examples are Vietoris topology [34, 35, 36, 37] (see, also, [1, 2, 5, 7, 8, 4, 11, 12, 13, 9, 10, 24, 23, 14, 18]) and Fell topology [17]. In this article, we concentrate on an extension of Vietoris based on the strongly far proximity.

Vietoris topology

Let X be an Hausdorff space. The *Vietoris topology* on $\mathrm{CL}(X)$ has as subbase all sets of the form

- $V^- = \{E \in \mathrm{CL}(X) : E \cap V \neq \varnothing\}$, where V is an open subset of X,
- $W^+ = \{C \in \mathrm{CL}(X) : C \subset W\}$, where W is an open subset of X.

The topology $\tau_V{}^-$ generated by the sets of the first form is called **hit part** because, in some sense, the closed sets in this family hit the open sets V. Instead, the topology $\tau_V{}^+$ generated by the sets of the second form is called **miss part**, because the closed sets here miss the closed sets of the form $X \setminus W$.

The Vietoris topology is the join of the two part: $\tau_V = \tau_V{}^- \vee \tau_V{}^+$. It represents the prototype of hit and miss topologies.

The Vietoris topology was modified by Fell. He left the hit part unchanged and in the miss part, ${\tau_F}^+$ instead of taking all open sets W, he took only open subsets with compact complement.
Fell topology: $\tau_F = {\tau_V}^- \vee {\tau_F}^+$.

It is possible to consider several generalizations. For example, instead of taking open subsets with compact complement, for the miss part we can look at subsets running in a family of closed sets $\mathscr{B}$. So we define the *hit and miss topology on* $CL(X)$ *associated with* $\mathscr{B}$ as the topology generated by the join of the hit sets A^-, where A runs over all open subsets of X, with the miss sets A^+, where A is once again an open subset of X, but more, whose complement runs in $\mathscr{B}$.

Another kind of generalization concerns the substitution of the inclusion present in the miss part with a strong inclusion associated with a proximity. Namely, when the space X carries a proximity δ, then a proximity variation of the miss part can be displayed by replacing the miss sets with *far-miss sets* $A^{++} := \{ E \in \mathrm{CL}(X) : E \ll_\delta A \}$.

Also, in this case we can consider A with the complement running in a family $\mathscr{B}$ of closed subsets of X. Then the *hit and far-miss topology* , $\tau_{\delta,\mathscr{B}}$, *associated with* $\mathscr{B}$ is generated by the join of the hit sets A^-, where A is open, with far-miss sets A^{++}, where the complement of A is in $\mathscr{B}$.

Fell topology can be considered as well as an example of hit and far-miss topology. In fact, in any EF-proximity, when a compact set is contained in an open set, it is also strongly contained.

3. Main Results

When we consider a proximity δ, it can happen that $A\delta B$ and they have empty intersection or, if X is a topological space, there could be only one point in the intersection of their closures. We need something more. We want to talk about *strong nearness* [30] (see, also, [29, 33, 32]) for subsets that at least have some points in common. For this reason, we introduce a new kind of proximity, that we call *almost proximity.*



Fig. **3.1.** $\boldsymbol{A \overset{\wedge\wedge\wedge}{\delta} B,\ D \overset{\wedge\wedge\wedge}{\delta} E}$

Definition 3.1. *Let* X *be a topological space. We say that the relation* $\overset{\wedge\wedge\wedge}{\delta}$ *on* $\mathscr{P}(X)$ *is a strongly near proximity called an* almost proximity*, provided it satisfies the following axioms. Let* $A, B, C \subset X$ *and* $x \in X$*:*

N0) $\varnothing \overset{\wedge\wedge\wedge}{\not\delta} A, \forall A \subset X$, *and* $X \overset{\wedge\wedge\wedge}{\delta} A, \forall A \subset X$

N1) $A \overset{\wedge\wedge\wedge}{\delta} B \Leftrightarrow B \overset{\wedge\wedge\wedge}{\delta} A$

N2) $A \overset{\wedge\wedge\wedge}{\delta} B \Rightarrow A \cap B \neq \varnothing$

N3) *If* $int(B)$ *and* $int(C)$ *are not equal to the empty set,* $A \overset{\wedge\wedge\wedge}{\delta} B$ *or* $A \overset{\wedge\wedge\wedge}{\delta} C \Rightarrow A \overset{\wedge\wedge\wedge}{\delta} (B \cup C)$

N4) $intA \cap intB \neq \varnothing \Rightarrow A \overset{\wedge\wedge\wedge}{\delta} B$ ■

When we write $A \overset{\wedge\wedge\wedge}{\delta} B$, we read A is *strongly near* B.

Example 3.2. A simple example of *almost proximity* that is also a *Lodato proximity* for nonempty sets $A, B \subset X$ is $A \overset{\wedge\!\!\wedge}{\delta} B \Leftrightarrow A \cap B \neq \emptyset$. ■

Example 3.3. Another example of *almost proximity* is given for nonempty sets $A, B \subset X$ in Fig. 3.1, where $A \overset{\wedge\!\!\wedge}{\delta} B \Leftrightarrow \text{int}(A) \cap \text{ int}(B) \neq \emptyset$. This is not a usual proximity. ■

Example 3.4. We can define another *almost proximity* for nonempty sets $D, E \subset X$ in Fig. 3.1 in the following way: $D \overset{\wedge\!\!\wedge}{\delta} D \Leftrightarrow E \cap \text{ int}(D) \neq \emptyset$ or $\text{int}(E) \cap D \neq \emptyset$. ■

Furthermore, for each *almost proximity*, we assume the following relation:

N5) $x \in \text{int}(A) \Rightarrow \{x\} \overset{\wedge\!\!\wedge}{\delta} A$
N6) $\{x\} \overset{\wedge\!\!\wedge}{\delta} \{y\} \Leftrightarrow x = y$ ■

So, for example, if we take the almost proximity of example 3.3, we have that

$$A \overset{\wedge\!\!\wedge}{\delta} B \Leftrightarrow \text{int}A \cap \text{int}B \neq \emptyset,$$

provided A and B are not singletons and if $A = \{x\}$, then $x \in \text{int}(B)$ and if B is also a singleton, then $x = y$. It turns out that if $A \subset X$ is an open set, then each point that belongs to A is strongly near A.

Now we want to define a new kind of *hit and far-miss topology* on $\text{CL}(X)$ by using *almost proximities*. That is, let τ_δ be the hit and far-miss topology having as subbase the sets of the form:

- $V^- = \{E \in \text{CL}(X) : E \cap V \neq \emptyset\}$, where V is an open subset of X,
- $A^{++} = \{\ E \in \text{CL}(X) : E \not\delta\, X \setminus A\ \}$, where A is an open subset of X.

where δ is a Lodato proximity compatible with the topology on X, and $\tau^{\wedge\!\!\wedge}$ the *strongly hit and far-miss* topology having as subbase the sets of the form:

- $V^{\wedge\!\!\wedge} = \{E \in \text{CL}(X) : E \overset{\wedge\!\!\wedge}{\delta} V\}$, where V is an open subset of X,
- $A^{++} = \{\ E \in \text{CL}(X) : E \not\delta\, X \setminus A\ \}$, where A is an open subset of X.

It is possible to define *strongly hit and miss* topology in many other ways by simply mixing strongly hit sets with miss sets or also strongly miss sets in a manner similar to that given in [31].

Theorem 3.5. *If the space X is T_1, the* strongly hit and far-miss *topology is an admissible topology.*

Proof. To achieve the result we need a T_1 space X in order to work with closed singletons. The result about the far-miss part is already known. So look at the strongly hit part. We want that the canonical injection $i : X \to \text{CL}(X)$, defined by $i(x) = \{x\}$, is a homeomorphism. It is continuous because $i^{-1}(V^{\wedge\!\!\wedge}) = \{x \in X : x \overset{\wedge\!\!\wedge}{\delta} V\} = V$. Moreover it is open in the relative topology on $i(X)$ because, for each open set A, $i(A) = A^{\wedge\!\!\wedge} \cap i(X)$. □

Remark 3.6. *Observe that the usual hit topology is a special strongly hit topology when taking as* almost proximity *either the one in example 3.2 or the one in example 3.4.* ■

Now we consider comparisons between usual *hit and far miss topologies* and *strongly hit and far miss* ones. To this purpose we need the following lemma.

Lemma 3.7. *Let A and H be open subsets of a T_1 topological space X. Then $A^- \subseteq H^{\doublewedge}$ implies $A \subseteq H$.*

Proof. By contradiction, suppose $A \nsubseteq H$. Then there exists $a \in A$ such that $a \notin H$. Hence $\{a\} \cap A \neq \varnothing$ while $a \overset{\doublewedge}{\not\delta} H$ by property $(N2)$. This is absurd. $\square$

Now let $\tau_\delta{}^-$ be the hypertopology having as subbase the sets of the form V^-, where V is an open subset of X, and let $(\tau^{\doublewedge})^-$ the hypertopology having as subbase the sets of the form $V^{\doublewedge}$, again with V an open subset of X.

Theorem 3.8. *Let X be a T_1 topological space. The hypertopologies $\tau_\delta{}^-$ and $(\tau^{\doublewedge})^-$ are not comparable.*



Fig. **3.2.** $(C = B(O,1) \cup S^1(O,s)) \notin H^{\doublewedge}$

Proof. First, we prove that $\tau_\delta{}^- \notin (\tau^{\doublewedge})^-$. Take $A^- \in \tau_\delta{}^-$ and $E \in A^-$, with $E \in \mathrm{CL}(X)$. We ask whether there exists an open set H such that $E \in H^{\doublewedge} \subset A^-$. To this purpose, consider the almost proximity of example 3.3 and let E be such that $\mathrm{int}(E) = \varnothing$. Then $E \cap A \neq \varnothing$, but $\mathrm{int}(E) \cap H = \varnothing$ for each $H \subset X$.

Conversely, we want to prove that $(\tau^{\doublewedge})^- \notin \tau_\delta{}^-$. Suppose $X = \mathbb{R}^2$ and consider, as before, the almost proximity of example 3.3. Take the closed set $E = B(O,2)$, that is the closed ball with the origin as center and radius 2, and the open set H as in Fig. 2.2. So $E \overset{\doublewedge}{\delta} H$. By lemma 3.7, to identify an open set A such that $E \in A^- \subseteq H^{\doublewedge}$, we need to consider $A \subseteq H$. But now it is possible to choose $C \in A^-$ such that $C \notin H^{\doublewedge}$, for each $A \subset H$. For example, we can take $C = B(O,1) \cup S^1(O,s)$, where $S^1(O,s)$ is the circumference with the origin as center and radius s for a suitable value of s (see, *e.g.*, Fig. 3.2).

$\square$

References

[1] G. Beer, *Topologies on Closed and Closed Convex Sets*, Mathematics and its Applicationsm, 268. Kluwer Academic Publishers Group, Dordrecht, (1993), xii+340 pp. ISBN: 0-7923-2531-1, MR1269778.

[2] G. Beer, R.K. Tamaki, *On hit-and-miss hyperspace topologies*, Comment. Math. Univ. Carolin. **34** (1993), no. 4, 717728, MR1263801.

[3] E. Čech, *Topological Spaces*, Publishing House of the Czechoslovak Academy of Sciences, Prague; Interscience Publishers John Wiley & Sons Ltd., London, UK, 1966, from seminar, Brno, 1936-1939; rev. ed. Z. Frolík, M. Katětov, 893 pp, MR0211373.

[4] A. Di Concilio, *Uniformities, hyperspaces, and normality*, Monatsh. Math. **107** (1989), no. 3, 303-308, MR1012462.

[5] A. Di Concilio, *Action on hyperspaces*, Topology Proc. **41** (2013), 85-98, MR2923735.

[6] A. Di Concilio, *Proximity: a powerful tool in extension theory, function spaces, hyperspaces, boolean algebra and point-set topology*, F. Mynard and E. Pearl(eds), Beyond Topology, Contemporary Mathematics Amer. Math. Soc.,**486**, (2009), 89-114.

[7] A. Di Concilio, S.A. Naimpally, *Proximal set-open topologies*, Boll. Unione Mat. Ital. Sez. B Artic Ric. Mat. **8** (2000), no. 1, 173-191, MR1755708.

[8] A. Di Concilio, *Proximal set-open topologies on partial maps*, Acta Math. Hungar. **88** (2000), no. 3, 227-237, MR1767801.

[9] G. Di Maio, S.A. Naimpally, *Hit-and-far-miss topologies*, Mat. Vesnik **60** (2008), no. 1, 59-78, MR2403272.

[10] G. Di Maio, E. Meccariello, S.A. Naimpally, *Hyperspace and function space are duals*, Questions Answers Gen. Topology **25** (2007), no. 1, 23-43, MR2319474.

[11] G. Di Maio, Lj.D.R. Kočinac, *Some covering properties of hyperspaces*, Topology Appl. **155** (2008), no. 17-18, 19591969, MR2457978.

[12] G. Di Maio, L. Holá, *On hit-and-miss topologies*, Rendiconto dell'Accademia delle Scienze Fisiche e Matematiche (4) **LXII** (1995), 103-124, MR1419286.

[13] G. Di Maio, S.A. Naimpally, *Comparison of hypertopologies*, Rend. Istit. Mat. Univ. Trieste **22** (1992), no. 1-2, 140-161, MR1210485.

[14] G. Di Maio, E. Meccariello, S.A. Naimpally, *Uniformly discrete hit-and-miss hypertopology. A missing link in hypertopologies*, Appl. Gen. Topol. **7** (2006), no. 2, 245-252, MR2295173.

[15] V.A. Efremovič, *The geometry of proximity* (Russian), Mat. Sbornik N. S. 31 (1952), 73, 189200, MR0055659.

[16] V.A. Efremovič, *Infinitesimal spaces* I (Russian), Doklady Akad. Nauk SSSR (N.S.) 76 (1951), 341343, MR0040748.

[17] J.M.G. Fell, *A Hausdorff topology for the closed subsets of a locally compact non-Hausdorff space*, Proc. Amer. Math. Soc. **13** (1962), 472-476, MR0139135.

[18] C. Guadagni, *Bornological Convergences on Local Proximity Spaces and ω_μ-Metric Spaces*, Ph.D thesis, Università degli Studi di Salerno, Dipartimento di Matematica, supervisor: A. Di Concilio, 2015, 72pp.

[19] C. Kuratowski, *Topologie. I et II. (French) [Topology. I and II]*, Éditions Jacques Gabay, Sceaux, 1992. iv+266 pp. ISBN: 2-87647-141-8, MR1296876.

[20] M.W. Lodato, *On topologically induced generalized proximity relations*, Ph.D. thesis, Rutgers University, 1962, 42pp, MR2613676.

[21] M.W. Lodato, *On topologically induced generalized proximity relations* I, Proc. Amer. Math. Soc. **15** (1964), 417-422, MR0161305.

[22] M.W. Lodato, *On topologically induced generalized proximity relations* II, Pacific J. Math. **17** (1966), 131-135, MR0192470.

[23] R. Lucchetti, *Hypertopologies and applications. Recent developments in well-posed variational problems*, Math. Appl. **331** (1995), Kluwer Acad. Publ., Dordrecht, 193-209, MR1351745.

[24] R. Lucchetti, A. Pasquale, *A new approach to a hyperspace theory*, J. of Convex Analysis **1** (1994), no. 2, 173-193, MR1363110.

[25] S.A. Naimpally, B.D. Warrack, *Proximity spaces*, Cambridge University Press, Cambridge Tract in Mathematics and Mathematical Physics 59, Cambridge, UK, 1970, ISBN 978-0-521-09183-1, MR2573941.

[26] S.A. Naimpally, *All hypertopologies are hit-and-miss*, App. Gen. Topology **3** (2002), 197-199.

[27] S.A. Naimpally, *Proximity Approach to Problems in Topology and Analysis*, Oldenbourg Verlag, Munich, Germany, 2009, 73 pp., ISBN 978-3-486-58917-7, MR2526304.

[28] S.A. Naimpally, J.F. Peters, *Topology with applications. Topological spaces via near and far. With a foreword by Iskander A. Taimanov.* World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013. xvi+277 pp. ISBN: 978-981-4407-65-6, MR3075111.

[29] J.F. Peters, *Proximal Voronoï regions, convex polygons, & Leader uniform topology*, Advances in Math. **4** (2015), no. 1, 1-5.

[30] J.F. Peters, *Visibility in proximal Delaunay meshes*, Advances in Math. (2015), *to appear.*

[31] J.F. Peters, C. Guadagni, *Strongly far proximity and hyperspace topology*, arXiv[Math.GN] **1502** (2015), no. 02771, 1-6.

[32] J.F. Peters, S.A. Naimpally, *Applications of near sets*, Notices Amer. Math. Soc. **59** (2012), no. 4, 536-542, MR2951956.

[33] J.F. Peters, M.A. Öztürk, M. Uçkun, *Klee-Phelps convex groupoids*, arXiv [math.GR] **1411** (2014), no. 0934, 1-4.

[34] L. Vietoris, *Stetige mengen*, Monatshefte fr Mathematik und Physik **31** (1921), no. 1, 173-204, MR1549101.

[35] L. Vietoris, *Bereiche zweiter ordnung*, Monatshefte fr Mathematik und Physik **32** (1922), no. 1, 258-280, MR1549179.

[36] L. Vietoris, *Kontinua zweiter ordnung*, Monatshefte fr Mathematik und Physik **33** (1923), no. 1, 49-62, MR1549268.

[37] L. Vietoris, *Über den höheren Zusammenhang kompakter Räume und eine Klasse von zusammenhangstreuen Abbildungen*, Math. Ann. **97** (1927), no. 1, 454-472, MR1512371.

[38] S. Willard, *General topology,* Addison-Wesley, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. (1970), xii+369 pp., MR0264581.

E-mail address: `James.Peters3@umanitoba.ca, cguadagni@unisa.it`

[α] Computational Intelligence Laboratory, University of Manitoba, WPG, MB, R3T 5V6, Canada and Department of Mathematics, Faculty of Arts and Sciences, Adiyaman University, 02040 Adiyaman, Turkey

[β] Computational Intelligence Laboratory, University of Manitoba, WPG, MB, R3T 5V6, Canada and Department of Mathematics, University of Salerno, via Giovanni Paolo II 132, 84084 Fisciano, Salerno , Italy